


BJ



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 0 891 975 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:
20.01.1999 Bulletin 1999/03

(51) Int. Cl.⁶: C07D 401/12, C07D 401/14,
C07D 405/14, C07D 417/14,
A01N 43/56, A01N 43/78

(21) Application number: 97914601.6

(22) Date of filing: 04.04.1997

(86) International application number:
PCT/JP97/01162

(87) International publication number:
WO 97/37990 (16.10.1997 Gazette 1997/44)

(84) Designated Contracting States:
DE DK ES FR GB IE IT NL PT

(30) Priority: 05.04.1996 JP 83587/96
09.08.1996 JP 211377/96

(71) Applicant:
MITSUBISHI CHEMICAL CORPORATION
Chiyoda-ku, Tokyo 100-0005 (JP)


(72) Inventors:
• KYOMURA, Nobuo
Aoba-ku, Yokohama-shi, Kanagawa 227 (JP)
• IKEDA, Yoshiya
Aoba-ku, Yokohama-shi, Kanagawa 227 (JP)

• OKUI, Shuko
Aoba-ku, Yokohama-shi, Kanagawa 227 (JP)
• TOMITA, Hirofumi
Aoba-ku, Yokohama-shi, Kanagawa 227 (JP)
• HIGASHINO, Yoshiaki
Aoba-ku, Yokohama-shi, Kanagawa 227 (JP)
• KOIKE, Sanae
Aoba-ku, Yokohama-shi, Kanagawa 227 (JP)

(74) Representative:
VOSSIUS & PARTNER
Siebertstrasse 4
81675 München (DE)

(54) PYRAZOLES AND AGRICULTURAL CHEMICALS CONTAINING THEM AS ACTIVE
INGREDIENTS

(57) A pyrazole compound represented by formula (I):




(I)

wherein R¹ represents a hydrogen atom, a C₁₋₄ alkyl group, a C₁₋₄ haloalkyl group, a C₁₋₄ alkoxy group or a phenyl group; R² represents a hydrogen atom, a halogen atom, a hydroxyl group, a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, a C₁₋₄ haloalkoxy group, a C₂₋₈ alkoxyalkoxy group or a benzyloxy group; or R₁ and R₂ are taken together to form a group:



wherein R^4 represents a hydrogen atom or a C_{1-4} alkyl group;

R^3 represents a hydrogen atom, a C_{1-4} alkyl group, a C_{2-5} alkenyl group, a C_{5-7} cycloalkenyl group, a C_{2-5} alkoxyalkyl group, a C_{8-12} benzyloxyalkyl group, a formyl group, a C_{2-5} alkanoyl group, a benzoyl group, a C_{2-5} alkoxy carbonyl group, a benzyloxycarbonyl group, a C_{1-4} alkylsulfonyl group or a C_{6-10} arylsulfonyl group; and R represents a group represented by formulae:



wherein R^5 represents a hydrogen atom, a halogen atom or a C_{1-4} alkoxy group; R^6 represents a halogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a C_{1-4} haloalkoxy group; and n represents 1 or 2, and a pesticide, such as a fungicide, a bacteriocide, an insecticide or an acaricide, comprising the same as an active ingredient. The pyrazole compound exhibits higher pesticidal activities than conventional pesticidal compounds and are of high safety.

Description

Technical Field:

- 5 This invention relates to a novel pyrazole compound and a pesticide (e.g., fungicides, bacteriocides, insecticides and acaricides) containing the same.

Background Art:

- 10 In agriculture and horticulture, a wide variety of pesticides such as fungicides, bacteriocides, insecticides and acaricides have been developed and put to practical use for the purpose of controlling pests.

However, currently spread agricultural and horticultural fungicides, bacteriocides, insecticides and acaricides are not necessarily satisfactory in pesticidal effect, pesticidal spectrum, residual activity, and the like. Further they may be seen as inadequate to meet the demand for reducing the times of application or the amount to be applied.

- 15 On the other hand, emergence of strains of pathogenic microorganisms or insect pests which have acquired resistance to the conventional pesticides has now become a serious problem. For example, pathogenic microorganisms or insect pests having developed resistance to various types of pesticides, such as triazole compounds, imidazole compounds, pyrimidine compounds, benzimidazole compounds, dicarboxyimide compounds, phenylamide compounds and organophosphorus compounds, have come out in various sites of cultivation of vegetables, fruits, flowers, teas, grains, 20 rices, etc. Control of various plant diseases or damages caused by these pathogens or insect pests is becoming increasingly difficult year by year.

While there are some pesticides to which pathogens or insect pests have not yet acquired resistance, such as dithiocarbamate compounds and phthalimide compounds, they are not preferred from the viewpoint of environmental pollution because, in general, they should be applied in large amounts or frequently.

- 25 Accordingly, it has been keenly demanded to develop a novel fungicide, bactericide or insecticide which exhibits sufficient controlling effects at a low amount on various pathogenic fungi or bacteria and insect pests having acquired resistance to the conventional agricultural or horticultural fungicides, bacteriocides and insecticides with minimal influences on the environment. As for acaricides, too, it has been expected to develop an acaricide which exhibits an excellent controlling effect on mites having resistance against conventional acaricides while securing safety.

- 30 JP-A-7-173139 (the term "JP-A" as used herein means an "unexamined published Japanese patent application") discloses N-(substituted pyridylmethyl)pyrazolecarboxamide compounds, teaching that the compounds have fungicidal, bactericidal, insecticidal and acaricidal activities. However, the substituents on the pyridine ring disclosed in the publication are limited to acyclic groups.

- 35 N-(Phenoxy pyridylmethyl)pyrazolecarboxamide compounds are proposed in JP-A-2-62876 as compounds having fungicidal, bactericidal, insecticidal and acaricidal activities. However, almost all the compounds disclosed are those having a substituent on the 4-position of the phenoxy moiety, and there is presented only one 3,4-disubstituted compound as a compound having a substituent on the 3-position of the phenoxy moiety.


- An object of the present invention is to provide a chemical substance which exhibits high controlling effects on various pathogens and insect pests showing resistance to the conventional agricultural or horticultural pesticides and is 40 useful as an active ingredient of pesticides such as fungicides, bacteriocides, insecticides and acaricides. Another object of the present invention is to provide highly safe pesticides which have the above-mentioned characteristics and yet are less involved in the problems of residual toxicity and environmental pollution.

- In order to achieve the above objects, the inventors of the present invention have carried out extensive studies and, as a result, found that a pyrazole compound represented by the formula shown below which possesses (1) a phenoxy 45 group having a specific condensed ring or (2) a phenoxy group having a specific substituent on the meta-position thereof bears all the above-described characteristics. The present invention has been completed based on this finding.

Disclosure of the Invention:


The gist of the present invention consists in a pyrazole compound represented by formula (I) and a pesticide comprising the compound as an active ingredient:

50




(I)

wherein R^1 represents a hydrogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a phenyl group; R^2 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_{1-4} alkyl group, a C_{1-4} alkoxy group, a C_{1-4} haloalkoxy group, a C_{2-8} alkoxyalkoxy group or a benzyloxy group; or R_1 and R_2 are taken together to form a group:



wherein R^4 represents a hydrogen atom or a C_{1-4} alkyl group;

R^3 represents a hydrogen atom, a C_{1-4} alkyl group, a C_{2-5} alkenyl group, a C_{5-7} cycloalkenyl group, a C_{2-5} alkoxyalkyl group, a C_{8-12} benzyloxyalkyl group, a formyl group, a C_{2-5} alkanoyl group, a benzoyl group, a C_{2-5} alkoxycarbonyl group, a benzyloxycarbonyl group, a C_{1-4} alkylsulfonyl group or a C_{6-10} arylsulfonyl group; and R represents a group represented by formulae:



wherein R^5 represents a hydrogen atom, a halogen atom or a C_{1-4} alkoxy group; R^6 represents a halogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a C_{1-4} haloalkoxy group; and n represents 1 or 2.

Hereinafter, the present invention is explained in detail.

The pyrazole compounds according to the present invention are compounds represented by formula (I), wherein:

R^1 represents a hydrogen atom; a C_{1-4} straight-chain or branched alkyl group, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl; a C_{1-4} straight-chain or branched haloalkyl group, such as difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, 3-chloropropyl, 3-bromopropyl, 3,3,3-trifluoropropyl, 3,3,3-trichloropropyl, 2,2,3,3-tetrafluoropropyl, 2,2,3,3,3-pentafluoropropyl group, 2,2-dichloro-3,3,3-trifluoropropyl, 1,3-difluoro-2-propyl, 1,1,1,3,3,3-hexafluoro-2-propyl, 4-chlorobutyl, 4,4,4-trifluorobutyl or 3,3,4,4,4-pentafluorobutyl; a C_{1-4} straight-chain or branched alkoxy group, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or t-butoxy; or a phenyl group.

R^2 represents a hydrogen atom; a halogen atom, such as fluorine, chlorine or iodine; a hydroxyl group; a C_{1-4} straight-chain or branched alkyl group, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl; a C_{1-4} straight-chain or branched alkoxy group, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or t-butoxy; a C_{1-4} straight-chain or branched haloalkoxy group, such as difluoromethoxy, trifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2,2,2-trifluoroethoxy, 2,2,2-trichloroethoxy, 3-chloropropoxy, 3-bromopropoxy, 3,3,3-trifluoropropoxy, 2,2,3,3-tetrafluoropropoxy, 2,2,3,3,3-pentafluoropropoxy, 2,2-dichloro-3,3,3-trifluoropropoxy, 1,3-difluoro-2-propoxy, 1,1,1,3,3,3-hexafluoro-2-propoxy, 3,3,3-trichloropropoxy, 4-chlorobutoxy, 4,4,4-trifluorobutoxy or 3,3,4,4,4-pentafluorobutoxy; or a benzyloxy group.

R¹ and R² may be taken together to form a group represented by formula:



wherein R⁴ represents a hydrogen atom or a C₁₋₄ straight-chain or branched alkyl group, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl. Of the groups formed by R¹ and R² connected together, a group represented by formula:



is preferred, in which R⁴ is at the 6-position of the condensed pyrazole ring.

R³ represents a hydrogen atom; a C₁₋₄ straight-chain or branched alkyl group, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl; a C₂₋₅ straight-chain or branched alkenyl group, such as vinyl, propenyl, butenyl or pentenyl; a C₅₋₇ cycloalkenyl group, such as cyclopentenyl, cyclohexenyl or cycloheptenyl; a C₂₋₅ straight-chain or branched alkoxyalkyl group, such as methoxymethyl, ethoxymethyl, isopropoxymethyl, methoxyethyl, ethoxyethyl, methoxypropyl or ethoxypropyl; a C₈₋₁₂ benzyloxyalkyl group, such as benzyloxymethyl, benzyloxyethyl or benzyloxypropyl; a formyl group; a C₂₋₅ alkanoyl group, such as acetyl, propionyl, n-butyryl, n-valeryl, isovaleryl, 2-methylbutyryl or pivaloyl; a benzoyl group; a C₂₋₅ alkoxy carbonyl group, such as methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl or t-butoxycarbonyl; a benzyloxy carbonyl group; a C₁₋₄ alkylsulfonyl group, such as methanesulfonyl or ethanesulfonyl; or a C₆₋₁₀ arylsulfonyl group, such as p-toluenesulfonyl. R³ is preferably a hydrogen atom, a C₂₋₅ alkenyl group, a C₅₋₇ cycloalkenyl group, a C₂₋₅ alkoxyalkyl group, a benzyloxymethyl group, a formyl group, a C₂₋₅ alkanoyl group or a p-toluenesulfonyl group, with a hydrogen atom, a formyl group or an alkanoyl group being particularly preferred.

R⁵ represents a hydrogen atom; a halogen atom, such as fluorine, chlorine or iodine; or a C₁₋₄ straight-chain or branched alkoxy group, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or t-butoxy.


R⁶ represents a halogen atom, such as fluorine, chlorine or iodine; a C₁₋₄ straight-chain or branched alkyl group, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl or t-butyl; a C₁₋₄ straight-chain or branched haloalkyl group, such as difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, 3-chloropropyl, 3-bromopropyl, 3,3,3-trifluoropropyl, 2,2,3,3-tetrafluoropropyl, 2,2,3,3,3-pentafluoropropyl group, 2,2-dichloro-3,3,3-trifluoropropyl, 1,3-difluoro-2-propyl, 1,1,1,3,3,3-hexafluoro-2-propyl, 3,3,3-trichloropropyl, 4-chlorobutyl, 4,4,4-trifluorobutyl or 3,3,4,4,4-pentafluorobutyl; a C₁₋₄ straight-chain or branched alkoxy group, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or t-butoxy; or a C₁₋₄ straight-chain or branched haloalkoxy group, such as difluoromethoxy, trifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2,2,2-trifluoroethoxy, 2,2,2-trichloroethoxy, 3-chloropropoxy, 3-bromopropoxy, 3,3,3-trifluoropropoxy, 2,2,3,3-tetrafluoropropoxy, 2,2,3,3,3-pentafluoropropoxy, 2,2-dichloro-3,3,3-trifluoropropoxy, 1,3-difluoro-2-propoxy, 1,1,1,3,3,3-hexafluoro-2-propoxy, 3,3,3-trichloropropoxy, 4-chlorobutoxy, 4,4,4-trifluorobutoxy or 3,3,4,4,4-pentafluorobutoxy.

R⁶ is preferably a halogen atom, a C₁₋₄ alkyl group, a C₁₋₄ haloalkyl group, a C₁₋₄ alkoxy group or a C₁₋₄ haloalkoxy group, with a halogen atom or a C₁₋₄ haloalkyl group being particularly preferred.

Where R is an meta-substituted phenoxy group, R¹ is preferably a C₁₋₄ alkyl group or a C₁₋₄ haloalkyl group, and R² is preferably a hydrogen atom, a halogen atom, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group.

Where R is a condensed phenoxy group, R¹ is preferably a hydrogen atom or a C₁₋₄ alkyl group, and R² is preferably a halogen atom, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group.

The compounds represented by formula (I) are prepared, for example, in accordance with the following reaction formula:



wherein R^1 , R^2 , R^3 , and R are as defined above; and Z represents a chlorine atom, a bromine atom, a hydroxyl group or a lower alkoxy group, such as methoxy, ethoxy or propoxy.

The compound (III) is used in an amount of 0.5 to 2.0 mol, preferably 0.8 to 1.5 mol, per mole of the compound (II).

Where Z in formula (II) is a chlorine atom or a bromine atom, the above reaction is carried out at 0° to 30°C , preferably 0° to 5°C , for 1 to 3 hours in a solvent in the presence of a base. Suitable solvents include aromatic hydrocarbons such as benzene, toluene and xylene; ketones such as acetone, methyl ethyl ketone and methyl isobutyl ketone; halogenated hydrocarbons such as chloroform and methylene chloride; water; esters such as methyl acetate and ethyl acetate; and polar solvents such as tetrahydrofuran, acetonitrile, dioxane, *N,N*-dimethylformamide, *N*-methylpyrrolidone, and dimethyl sulfoxide. Preferred of them are aromatic hydrocarbons, esters, tetrahydrofuran, acetonitrile, and dioxane. The solvent is used in an amount 3 to 30 times, preferably 5 to 20 times, the weight of the compound (III). Suitable bases include commonly employed inorganic bases, such as alkali metal hydroxides (e.g., sodium hydroxide and potassium hydroxide), alkaline earth metal hydroxides (e.g., magnesium hydroxide and calcium hydroxide), alkali metal oxides, and alkali metal carbonates; and organic bases, such as triethylamine, pyridine, 1,4-diazabicyclo[2.2.2]octane, 1,5-diazabicyclo[4.3.0]non-5-ene, and 1,8-diazabicyclo[5.4.0]undec-7-ene. An alkali metal hydroxide, triethylamine or pyridine is preferred. The base is used in an amount of 0.5 to 2.0 mol, preferably 0.8 to 1.5 mol, per mole of the compound (II).

Where Z in formula (II) is a hydroxyl group, the reaction is conducted at 150° to 250°C , preferably 200° to 250°C , in a solvent under atmospheric pressure or under pressure. Suitable solvents include high-boiling solvents such as *N,N*-dimethylformamide, *N*-methylpyrrolidone, and dimethyl sulfoxide. Where Z in formula (II) is a lower alkoxy group, the reaction is performed at 150° to 250°C , preferably 200° to 250°C , with or without a solvent under atmospheric pressure or under pressure. Suitable solvents, if used, include high-boiling aprotic polar solvents such as *N,N*-dimethylformamide, *N*-methylpyrrolidone, and dimethyl sulfoxide.

The amount of the solvent, if used, is 3 to 30 times, preferably 5 to 20 times, the weight of the compound (III).

The compound (II) is prepared, for example, in accordance with the process described in *Bull. Soc. Chim. France*, 293 (1966) or *Annalen der Chemie, Justus Liebig's*, 536, 97 (1938). The compound (III) is prepared by, for example, the process described in *Journal fur Praktische Chemie*, 146, 95 (1936).

The compounds represented by formula (I) according to the present invention show high fungicidal effects on plant pathogenic fungi, such as blast fungi, rust fungi, and downy mildew fungi, and are useful as an active ingredient of agricultural or horticultural fungicides. The compounds of the present invention also have high controlling activities on larvae and imagoes of (1) *Lepidoptera*, such as *Spodoptera litura*, *Chilo suppressalis*, *Cuapha locrocis medinalis*, *Plutella xylostella*, and *Adoxophyes* sp., (2) *Hemiptera*, such as *Delphacidae* (e.g., *Sogatella furcifera*, *Nilaparvata lugens*, and *Laodelphax striatellus*), *Deltocephalidae* (e.g., *Nephotettix cincticeps*, *Empoasca onukii*, and *Tettigella viridis*), *Aphididae* (e.g., *Myzus persicae* and *Aphis gossypii*), *Aleyrodidae* (e.g., *Trialeurodes vaporariorum*), and *Pentahomidae* (e.g., *Plautia stali*), (3) *Coleoptera*, such as *Phyllotreta striolata*, *Aulacophora femoralis*, and *Callosobruchus chinensis*, (4) *Diptera*, such as *Musca domestica*, *Aedes aegypti*, and *Culex pipiens pallens*, and (5) *Orthoptera*, such as *Periplaneta americana*, and also on eggs, larvae and imagoes of *Acarina*, such as *Tetranychus urticae*, *Tetranychus cinnabarinus*, and *Panonychus citri*, and are useful as an active ingredient of agricultural and horticultural insecticides and acaricides. The plant pathogens and insects which are controllable by the compounds of the present invention are not limited to the above examples.

In using the compounds of the present invention as pesticides, they may be applied as such but are preferably formulated into pesticidal compositions together with appropriate adjuvants customarily employed in the art. While not limiting, the forms of the pesticides include emulsifiable concentrates, wettable powders, dusts, flowables, powders, granules, tablets, oils, sprays, and fumigants. These preparations can comprise one or more than one compounds of the present invention.

The adjuvants are used in the pesticides for the purpose of improvement of the pesticidal effect, stabilization of the

reduced pressure to give 101.3 g of crude ethyl 1,3-dimethyl-4-chloropyrazole-5-carboxylate.

REFERENCE EXAMPLE 3

In a 1 l flask, 20.8 g of 95% sodium hydroxide was dissolved in 320 ml of water. To the solution was added 91.9 g of ethyl 1,3-dimethyl-4-chloropyrazole-5-carboxylate prepared in Reference Example 2. The mixture was vigorously stirred on an oil bath at 90°C for 1 hour to conduct the reaction (reaction temperature: 83°C). After cooling to room temperature, the reaction mixture was neutralized with diluted hydrochloric acid prepared from 50 ml of concentrated hydrochloric acid and 200 ml of water. The precipitate thus formed was collected by filtration, washed with water, and concentrated under reduced pressure to give 72.2 g (yield: 92.0%) of crude 1,3-dimethyl-4-chloropyrazole-5-carboxylic acid.

REFERENCE EXAMPLE 4

In a 100 ml flask were charged 9.3 g of 1,3-dimethyl-4-chloropyrazole-5-carboxylic acid obtained in Reference Example 3, 30 ml of toluene, and 11.9 g of thionyl chloride, and the mixture was heated under reflux for 2 hours on an oil bath. Concentration of the reaction mixture gave 9.3 g (yield: 91%) of crude 1,3-dimethyl-4-chloropyrazole-5-carboxylic acid chloride.

REFERENCE EXAMPLE 5

To a solution of 2.9 g of β -naphthol in 15 ml of dimethylformamide was slowly added 0.8 g of 60% sodium hydride, followed by stirring at room temperature for 30 minutes. To the mixture was added slowly 2.77 g of 2-chloro-5-cyanopyridine, followed by stirring at 40°C for 1 hour. After cooling to room temperature, 50 ml of ethyl acetate and 50 ml of water were added thereto. The extracted organic layer was washed with a saturated sodium chloride aqueous solution and dried over anhydrous sodium sulfate. Any insoluble matter was separated by filtration, and the filtrate was concentrated. The residue was purified by column chromatography on silica gel (developing solvent: hexane/ethyl acetate=4/1) to give 4.14 g of 2-(β -naphthoxy) 5-cyanopyridine.

REFERENCE EXAMPLE 6

To a solution of 1.72 g of 2-(β -naphthoxy)-5-cyanopyridine obtained in Reference Example 5 in 50 ml of ethanol were added 1.7 g of 28% aqueous ammonia and 1.5 g of Raney nickel to perform hydrogenation. After completion of the reaction, Raney nickel was separated by filtration, and the solvent was evaporated from the filtrate to recover crude 5-aminomethyl-2-(β -naphthoxy)pyridine.

EXAMPLE 1


Preparation of N-[6-(3-Trifluoromethylphenoxy)-3-pyridylmethyl]-1,3,4-trimethyl-5-pyrazolecarboxamide



A mixture of 7.7 g of 1,3,4-trimethyl-5-pyrazolecarboxylic acid and 12 g of thionyl chloride was heated under reflux for 1 hour. Thionyl chloride was removed by evaporation under reduced pressure, and the residue was dissolved in 20 ml of toluene. The resulting solution was added dropwise to a solution of 13.4 g of 5-aminomethyl-2-(3-trifluoromethylphenoxy)pyridine and 6.1 g of triethylamine in 30 ml of toluene at 0 to 10°C. After the addition, the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into ice-water and extracted with toluene. The toluene layer was washed successively with a sodium hydrogencarbonate aqueous solution, water and a saturated sodium chloride aqueous solution and dried over anhydrous sodium sulfate. The solvent was removed by evaporation under reduced pressure, and the residue was purified by silica gel column chromatography to give 18.2 g of compound No. 10 shown in Table 1 below.

EXAMPLE 2

The compounds shown in Table 1 were obtained in the same manner as in Example 1. In Table 1, the values with an asterisk in the column "Melting Point" are refractive indices.

TABLE 1



Compound No.	R ¹	R ²	R ⁵	M. P. (°C)
1	H	H	CF ₃	126-128
2	H	Cl	CF ₃	87-88
3	CH ₃	H	CF ₃	154.2-154.4 [†] (25°C)
4	CH ₃	Cl	CF ₃	118-119
5	CH ₃	Br	CF ₃	119-120
6	CH ₃	I	CF ₃	135-136
7	CH ₃	CH ₃	CH ₃	116-117
8	CH ₃	CH ₃	Cl	126-127
9	CH ₃	CH ₃	Br	118-120
10	CH ₃	CH ₃	CF ₃	125-126
11	CH ₃	CH ₃	OCHF ₂	106-107
12	C ₂ H ₅	H	CF ₃	155.1-155.8 [†] (25°C)
13	C ₂ H ₅	Cl	CF ₃	114-115
14	C ₂ H ₅	Br	CF ₃	127-128
15	C ₂ H ₅	CH ₃	CF ₃	121-122
16	iso-C ₃ H ₇	Cl	CF ₃	107-108
17	C ₆ H ₅	H	CF ₃	157.5-158.8 [†] (24°C)
18			CF ₃	119-120
19	H ₃ C- 		CF ₃	133-135
20	CF ₃	H	CF ₃	97-98

The NMR and IR spectra of Compound Nos. 3, 10, 12, and 17 were as follows.

Compound No. 3

¹H-NMR(CDCl₃) δ ppm: 2.24 (3H, s), 4.11 (3H, s), 4.54 (2H, d), 6.27 (1H, s), 6.37 (1H, bs), 6.96 (1H, d), 7.31 (1H, d), 7.39 (1H, s), 7.49 (2H, m), 7.75 (1H, dd), 8.12 (1H, d)
 IR(KBr) cm⁻¹: 3320, 2940, 1660

Compound No. 10

¹H-NMR(CDCl₃) δ ppm: 2.12 (3H, s), 2.18 (3H, s), 4.02 (3H, s), 4.60 (2H, d), 6.08 (1H, bs), 6.98 (1H, d), 7.31 (1H, d), 7.40 (1H, bs), 7.50 (2H, m), 7.78 (1H, dd), 8.16 (1H, d)
 IR(KBr) cm⁻¹: 3270, 1630, 1480, 1320, 1120, 920, 830

Compound No. 12

¹H-NMR(CDCl₃) δ ppm: 1.22 (3H, t), 2.62 (2H, q), 4.12 (3H, s), 4.55 (2H, d), 6.30 (1H, s), 6.33 (1H, bs), 6.97 (1H, d), 7.32 (1H, d), 7.40 (1H, s), 7.49 (2H, m), 7.75 (1H, dd), 8.13 (1H, d)
 IR(KBr) cm⁻¹: 3320, 2980, 1650, 1550, 1480, 1330, 1120, 920

Compound No. 17

¹H-NMR(CDCl₃) δ ppm: 4.24 (3H, s), 4.59 (2H, d), 6.38 (1H, bs), 6.77 (1H, s), 6.98 (1H, d), 7.4 (6H, m), 7.76 (3H, m), 8.18 (1H, s)
 IR(KBr) cm⁻¹: 3330, 2950, 1660, 1550, 1480, 1320, 1120, 920


EXAMPLE 3Preparation of N-[6-(β-Naphthyloxy)-3-pyridyl-methyl]-4-chloro-1,3-dimethyl-5-pyrazolecarboxamide

In 10 ml of toluene was dissolved 0.56 g of 4-chloro-1,3-dimethyl-5-pyrazolecarboxylic acid chloride. The resulting solution was added dropwise to a solution of 0.73 g of 5-aminomethyl-2-(β-naphthyloxy)pyridine and 0.3 g of triethylamine in 10 ml of toluene at 0 to 10°C. After the addition, the reaction mixture was stirred at room temperature for 2 hours, and then poured into ice-water and extracted with toluene. The toluene layer was washed successively with a sodium hydrogencarbonate aqueous solution, water, and a saturated sodium chloride aqueous solution and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the residue was purified by silica gel column chromatography to give 0.99 g of compound 22 (shown in Table 2).

EXAMPLE 4

The compounds shown in Tables 2 and 3 were obtained in accordance with the process of Example 3. In the Tables, the values with an asterisk are refractive indices.

TABLE 2



Compound No.	R ¹	R ²	R	M. Pt. (°C) or Refractive Index
2 1	CH ₃	Cl		1 1 6 - 1 1 7
2 2	CH ₃	Cl		1 5 0 - 1 5 1
2 3	CH ₃	Br		1 5 1 - 1 5 2
2 4	CH ₃	CH ₃		1 2 1 - 1 2 2
2 5	CH ₃	CH ₃		1 1 9 - 1 2 0
2 6	CH ₃	CH ₃		1 5 3 - 1 5 5
2 7	CH ₃	CH ₃		1 5 6 - 1 5 8

TABLE 2 (cont'd)






















Compound No.	R ¹	R ²	R	M. Pt. (°C) or Refractive Index
28	CH ₃	CH ₃		154 - 155
29	CH ₃	CH ₃		91 - 93
30	CH ₃	CH ₃		129 - 131
31	CH ₃	CH ₃		179 - 181
32	CH ₃	OH		amorphous solid
33	CH ₃	OCH ₃		85 - 87
34	CH ₃	OCH ₃		1.6109 [*] (24°C)
35	CH ₃	OCHF ₂		104 - 105
36	CH ₃			92 - 93
37	C ₂ H ₅	Cl		99 - 100
38	C ₂ H ₅	Cl		121 - 122
39	C ₂ H ₅	Br		127 - 129
40	C ₂ H ₅	CH ₃		126 - 128
41	C ₃ H ₇ ⁱ	Cl		122 - 123
42				157 - 158
43	H ₃ C- 			142 - 144

TABLE 3



Compound No.	R ¹	R ²	R ³	M. Pt. (°C) or Refractive Index
44	CH ₃	Cl	CH ₃	1.6038*(25°C)
45	C ₂ H ₅	Cl		82-83
46	C ₂ H ₅	Cl		viscous
47	C ₂ H ₅	Cl		viscous
48	CH ₃	OCH ₃	CHO	viscous
49	C ₂ H ₅	Cl	CHO	viscous
50	CH ₃	CH ₃	CHO	viscous
51	CH ₃	Cl	COCH ₃	1.6010*(25°C)
52	CH ₃	OCH ₃	COCH ₃	106
53	C ₂ H ₅	Cl	COCH ₃	viscous
54	CH ₃	CH ₃	COCH ₃	viscous
55	CH ₃	OCH ₃	COC ₂ H ₅	viscous
56	CH ₃	OCH ₃	COC ₃ H ₇	viscous
57	CH ₃	CH ₃	COC ₃ H ₇	viscous

TABLE 3 (cont'd)



Compound No.	R ¹	R ²	R ³	M. Pt. (°C) or Refractive Index
58	CH ₃	OCH ₃	COC ₆ H ₅	amorphous solid
59	CH ₃	OCH ₃	COOCH ₃	104-106
60	C ₂ H ₅	Cl	COOCH ₃	viscous
61	CH ₃	OCH ₃	COOCH ₂ C ₆ H ₅	amorphous solid
62	CH ₃	OCH ₃	CH ₂ OCH ₃	viscous
63	CH ₃	OCH ₃	CH ₂ OC ₂ H ₅	viscous
64	CH ₃	OCH ₃	CH ₂ OCH ₂ C ₆ H ₅	viscous
65	CH ₃	OCH ₃	SO ₂ CH ₃	viscous
66	CH ₃	OCH ₃	SO ₂ C ₆ H ₄ -p-CH ₃	viscous
67	C ₂ H ₅	Cl	CH ₃	143-144
68	C ₂ H ₅	Cl	COC ₂ H ₅	viscous
69	C ₂ H ₅	Cl	COC ₃ H ₇	viscous
70	C ₂ H ₅	Cl	COC ₆ H ₅	viscous
71	C ₂ H ₅	Cl	SO ₂ C ₆ H ₄ -p-CH ₃	viscous

The NMR data of the compounds shown in Tables 2 and 3 were as follows.

Compound 21:

¹H-NMR(CDCl₃) δ ppm: 2.11 (2H, m), 2.90 (4H, m), 4.13 (3H, s), 4.58 (2H, d), 6.89 (2H, d), 6.98 (1H, s), 7.00 (1H, bs), 7.22 (1H, d), 7.67 (1H, dd), 8.17 (1H, d)

Compound 22:

¹H-NMR(CDCl₃) δ ppm: 2.23 (3H, s), 4.12 (3H, s), 4.59 (2H, d), 6.96 (1H, d), 7.00-7.10 (1H, bt), 7.29 (1H, dd), 7.40-7.50 (2H, m), 7.56 (1H, d), 7.71-7.76 (2H, m), 7.86 (2H, t), 8.18 (1H, d)

Compound 23:

¹H-NMR(CDCl₃) δ ppm: 2.24 (3H, s), 4.13 (3H, s), 4.60 (2H, d), 6.97 (1H, d), 7.00-7.10 (1H, bt), 7.29 (1H, dd), 7.42-7.53 (2H, m), 7.56 (1H, d), 7.76 (2H, t), 7.86 (2H, t), 8.20 (1H, d)

Compound 24:

¹H-NMR(CDCl₃) δ ppm: 2.10 (2H, m), 2.10 (3H, s), 2.17 (3H, s), 2.90 (4H, m), 4.01 (3H, s), 4.57 (2H, d), 6.00 (1H, bs), 6.90 (2H, dd), 6.98 (1H, s), 7.22 (1H, d), 7.70 (1H, dd), 8.15 (1H, d)

Compound 25:

¹H-NMR(CDCl₃) δ ppm: 1.79 (4H, m), 2.10 (3H, s), 2.17 (3H, s), 2.76 (4H, bm), 4.01 (3H, s), 4.57 (2H, d), 6.00 (1H, bs), 6.85 (3H, m), 7.07 (1H, d), 7.68 (1H, dd), 8.17 (1H, s)

Compound 26:

¹H-NMR(CDCl₃) δ ppm: 2.11 (3H, s), 2.17 (3H, s), 4.01 (3H, s), 4.59 (2H, d), 6.05 (1H, bs), 6.97 (1H, d), 7.28 (1H, dd), 7.46 (2H, m), 7.56 (1H, d), 7.70-7.90 (4H, m), 8.17 (1H, d)

Compound 27:

¹H-NMR(CDCl₃) δ ppm: 2.12 (3H, s), 2.18 (3H, s), 4.02 (3H, s), 4.60 (2H, d), 6.05 (1H, bs), 6.99 (1H, d), 7.31 (1H, dd), 7.55 (2H, m), 7.64 (1H, d), 7.77 (2H, m), 8.01 (1H, s), 8.17 (1H, d)

Compound 28:

¹H-NMR(CDCl₃) δ ppm: 2.11 (3H, s), 2.18 (3H, s), 3.92 (3H, s), 4.01 (3H, s), 4.59 (2H, d), 6.02 (1H, bs), 6.95 (1H, d), 7.16 (2H, m), 7.3 (1H, d), 7.50 (1H, d), 7.70 (2H, m), 7.78 (1H, d), 8.17 (1H, d)

Compound 29:

¹H-NMR(CDCl₃) δ ppm: 1.34 (6H, s), 1.80 (2H, t), 2.10 (3H, s), 2.17 (3H, s), 2.77 (2H, t), 4.01 (3H, s), 4.57 (2H, d), 6.00 (1H, bs), 6.80 (3H, m), 7.68 (1H, dd), 8.17 (1H, d)

Compound 30:

¹H-NMR(CDCl₃) δ ppm: 2.12 (3H, s), 2.18 (3H, s), 4.02 (3H, s), 4.60 (2H, d), 6.10 (1H, b), 7.03 (1H, d), 7.40 (1H, m), 7.52 (2H, m), 7.78 (1H, dd), 8.13 (3H, m), 8.88 (1H, t)

Compound 31:

¹H-NMR(CDCl₃) δ ppm: 2.12 (3H, s), 2.18 (3H, s), 4.02 (3H, s), 4.60 (2H, d), 6.07 (1H, b), 6.99 (1H, d), 7.30 (1H, d), 7.74 (1H, dd), 7.78 (1H, dd), 8.15 (2H, d), 8.96 (1H, s)

Compound 32:

¹H-NMR(CDCl₃) δ ppm: 2.09 (3H, s), 4.03 (3H, s), 4.52 (2H, d), 6.88 (1H, d), 7.22 (1H, dd), 7.45 (2H, m), 7.49 (1H,

d), 7.64 (1H, m), 7.72 (2H, m), 7.83 (2H, m), 8.10 (1H, s)

Compound 33:

5 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.10 (2H, m), 2.28 (3H, s), 2.90 (4H, q), 3.81 (3H, s), 4.09 (3H, s), 4.54 (2H, d), 6.87 (2H, d), 6.98 (1H, bs), 7.22 (1H, d), 7.49 (1H, bs), 7.68 (1H, dd), 8.15 (1H, d)

Compound 34:

10 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.28 (3H, s), 3.82 (3H, s), 4.10 (3H, s), 4.57 (2H, d), 6.96 (1H, d), 7.28 (1H, dd), 7.46 (2H, m), 7.53 (1H, bs), 7.58 (1H, d), 7.70-7.90 (4H, m), 8.17 (1H, d)

Compound 35:

15 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.23 (3H, s), 4.11 (3H, s), 4.56 (2H, d), 6.35 (1H, t), 6.88 (1H, bs), 6.97 (1H, d), 7.28 (1H, dd), 7.46 (2H, m), 7.56 (1H, d), 7.70 (1H, d), 7.77 (1H, d), 7.86 (2H, t), 8.17 (1H, s)

Compound 36:

20 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.27 (3H, s), 4.10 (3H, s), 4.37 (2H, d), 4.93 (2H, s), 6.87 (1H, d), 7.21 (2H, m), 7.34 (3H, m), 7.46 (3H, m), 7.56 (2H, m), 7.78 (1H, d), 7.86 (2H, m), 8.04 (1H, s)

Compound 37:

25 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.23 (3H, t), 2.11 (2H, m), 2.63 (2H, q), 2.90 (4H, m), 4.13 (3H, s), 4.58 (2H, q), 6.89 (2H, d), 6.98 (1H, d), 7.00 (1H, bs), 7.22 (1H, d), 7.69 (1H, dd), 8.18 (1H, d)

Compound 38:

30 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.23 (3H, t), 2.63 (2H, q), 4.14 (3H, s), 4.60 (2H, d), 6.97 (1H, d), 7.00-7.20 (1H, bt), 7.27 (1H, dd), 7.40-7.50 (2H, m), 7.56 (1H, d), 7.76 (2H, m), 7.86 (2H, t), 8.19 (1H, d)

Compound 39:

35 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.23 (3H, t), 2.62 (2H, q), 4.14 (3H, s), 4.61 (2H, d), 6.97 (1H, d), 7.05 (1H, bs), 7.28 (1H, dd), 7.46 (2H, m), 7.55 (1H, d), 7.76 (2H, m), 7.86 (2H, t), 8.20 (1H, s)

Compound 40:

40 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.20 (3H, t), 2.13 (3H, s), 2.57 (2H, q), 4.02 (3H, s), 4.60 (2H, d), 6.06 (1H, bs), 6.97 (1H, d), 7.29 (1H, d), 7.46 (2H, m), 7.56 (1H, d), 7.77 (2H, m), 7.86 (2H, t), 8.18 (1H, d)

Compound 41:

45 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.28 (6H, d), 3.03 (1H, m), 4.13 (3H, s), 4.60 (2H, d), 6.97 (1H, d), 7.05 (1H, bs), 7.31 (1H, m), 7.46 (2H, m), 7.56 (1H, d), 7.76 (2H, m), 7.86 (2H, t), 8.19 (1H, d)

Compound 42:

50 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.46 (2H, m), 2.72 (4H, m), 4.15 (3H, s), 4.56 (2H, d), 6.02 (1H, bs), 6.96 (1H, d), 7.27 (1H, dd), 7.46 (2H, m), 7.55 (1H, d), 7.75 (2H, m), 7.86 (2H, t), 8.15 (1H, d)

Compound 43:

55 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.27 (3H, d), 2.00 (1H, m), 2.68 (3H, m), 3.12 (1H, m), 4.16 (3H, s), 4.56 (2H, d), 6.00 (1H, bs), 6.96 (1H, d), 7.28 (1H, dd), 7.46 (2H, m), 7.56 (1H, d), 7.76 (2H, m), 7.86 (2H, t), 8.15 (1H, d)

Compound 44:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.22 (3H, s), 2.98+3.03 (3H, s+s), 3.83+3.80 (3H, s+s), 4.71+4.52 (2H, s+b), 6.97 (1H, t), 7.30 (1H, dd), 7.47 (2H, m), 7.58 (1H, m), 7.78 (2H, m), 7.90 (2H, m), 8.18 (1H, d)

Compound 45:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.24 (3H, t), 2.63 (2H, q), 3.92 (3H, s), 4.99 (2H, s), 6.92 (1H, d), 7.28 (1H, dd), 7.45 (2H, m), 7.55 (1H, d), 7.75-7.83 (3H, m), 7.87 (1H, d), 7.29 (1H, d), 8.89 (1H, s)

Compound 46:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.23 (3H, s), 1.83 (2H, m), 2.19 (2H, bm), 2.30 (2H, bm), 3.81 (3H, s), 4.82 (2H, s), 5.14 (1H, bs), 6.95 (1H, d), 7.29 (1H, dd), 7.45 (2H, m), 7.57 (1H, d), 7.77 (2H, t), 7.86 (2H, t), 8.13 (1H, d)

Compound 47:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.26 (3H, s), 1.42 (2H, bm), 1.88 (2H, b), 1.99 (2H, b), 2.59 (2H, q), 3.78 (3H, s), 4.78 (2H, s), 5.29 (1H, bs), 6.95 (1H, d), 7.30 (1H, dd), 7.46 (2H, m), 7.57 (1H, d), 7.80 (2H, t), 7.86 (2H, t), 8.14 (1H, s)

Compound 48:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.25 (3H, s), 3.59 (3H, s), 3.91 (3H, s), 4.97 (2H, s), 6.90 (1H, d), 7.25 (1H, dd), 7.44 (2H, m), 7.53 (1H, d), 7.7-7.9 (4H, m), 8.30 (1H, d), 8.87 (1H, s)

Compound 49:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.24 (3H, t), 2.63 (2H, q), 3.92 (3H, s), 4.99 (2H, s), 6.92 (1H, d), 7.28 (1H, dd), 7.45 (2H, m), 7.55 (1H, d), 7.75-7.83 (3H, m), 7.87 (1H, d), 7.29 (1H, d), 8.89 (1H, s)

Compound 50:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.87 (3H, t), 2.18 (3H, s), 3.85 (3H, s), 4.98 (2H, s), 6.92 (1H, d), 7.27 (1H, dd), 7.46 (2H, m), 7.56 (1H, d), 7.76-7.89 (4H, m), 8.31 (1H, d), 8.86 (1H, s)

Compound 51:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.24 (3H, s), 2.28 (3H, s), 3.79 (3H, s), 4.95 (2H, b), 6.92 (1H, d), 7.26 (1H, m), 7.46 (2H, m), 7.55 (1H, d), 7.65 (1H, dd), 7.78 (1H, d), 7.86 (2H, m), 8.02 (1H, d)

Compound 52:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.23 (3H, s), 2.25 (3H, s), 3.66 (3H, s), 3.84 (3H, s), 4.98 (2H, s), 6.89 (1H, d), 7.24 (1H, d), 7.45 (2H, m), 7.53 (1H, d), 7.70 (1H, dd), 7.77 (1H, d), 7.84 (2H, m), 8.09 (1H, d)

Compound 53:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.24 (3H, t), 2.58 (3H, s), 2.62 (2H, q), 3.78 (3H, s), 4.80-5.10 (2H, b), 6.90 (1H, d), 7.25 (1H, dd), 7.44 (2H, m), 7.54 (1H, d), 7.64 (1H, dd), 7.76 (1H, dd), 7.82 (1H, dd), 7.85 (1H, d), 8.01 (1H, d)

Compound 54:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.94 (3H, t), 2.16 (3H, s), 2.18 (3H, s), 3.77 (3H, s), 4.8-5.1 (2H, b), 6.90 (1H, d), 7.26 (1H, dd), 7.46 (2H, m), 7.54 (1H, d), 7.67 (1H, d), 7.77 (1H, d), 7.84 (1H, dd), 7.87 (1H, d), 8.05 (1H, d)

Compound 55:

$^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.10 (3H, t), 2.44 (3H, s), 2.50 (2H, q), 3.65 (3H, s), 3.84 (3H, s), 4.97 (2H, s), 6.89 (1H,

d), 7.24 (1H, dd), 7.46 (2H, m), 7.52 (1H, d), 7.68-7.88 (4H, m), 8.09 (1H, d)

Compound 56:

5 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 0.84 (3H, s), 1.62 (2H, m), 2.42 (3H, s), 2.47 (2H, t), 3.66 (3H, s), 3.83 (3H, s), 4.96 (2H, s), 6.89 (1H, dd), 7.24 (1H, dd), 7.45 (2H, m), 7.52 (1H, d), 7.70 (1H, dd), 7.76 (1H, dd), 7.83 (1H, dd), 7.85 (1H, d), 8.09 (1H, d)

Compound 57:

10 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.24 (3H, t), 2.63 (2H, q), 3.92 (3H, s), 4.99 (2H, s), 6.92 (1H, d), 7.28 (1H, dd), 7.45 (2H, m), 7.55 (1H, d), 7.75-7.83 (3H, m), 7.87 (1H, d), 7.29 (1H, d), 8.89 (1H, s)

Compound 58:

15 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.93 (3H, s), 3.53 (3H, s), 3.75 (3H, s), 5.15 (2H, s), 6.93 (1H, d), 7.19-7.30 (4H, m), 7.39-7.50 (4H, m), 7.54 (1H, d), 7.74-7.88 (3H, m), 7.98 (1H, dd), 8.23 (1H, d)

Compound 59:

20 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.05 (3H, s), 3.59 (3H, s), 3.68 (3H, s), 3.93 (3H, s), 4.93 (2H, s), 6.92 (1H, d), 7.27 (1H, dd), 7.46 (2H, m), 7.55 (1H, d), 7.74-7.90 (3H, m), 8.30 (1H, d)

Compound 60:

25 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.22 (3H, t), 2.60 (2H, q), 3.69 (3H, s), 3.91 (3H, s), 4.94 (2H, bs), 6.93 (1H, d), 7.28 (1H, dd), 7.43 (2H, m), 7.56 (1H, d), 7.76 (1H, dd), 7.80-7.90 (3H, m), 8.28 (1H, d)

Compound 61:

30 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.15 (3H, s), 3.54 (3H, s), 3.88 (2H, s), 4.94 (2H, s), 5.09 (2H, s), 6.89 (1H, d), 7.06 (2H, m), 7.24-7.30 (4H, m), 7.46 (2H, m), 7.54 (1H, d), 7.74-7.85 (4H, m), 8.29 (1H, d)

Compound 62:

35 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.22 (3H, s), 3.01 (3H, s), 3.62 (3H, s), 3.81 (2H, s), 3.82 (2H, s), 4.72 (2H, s), 6.94 (1H, d), 7.29 (1H, dd), 7.46 (2H, m), 7.56 (1H, br), 7.77-7.89 (4H, m), 8.23 (1H, d)

Compound 63:

40 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 1.09 (3H, t), 2.22 (3H, s), 3.23 (2H, q), 3.63 (3H, s), 3.81 (2H, s), 4.74 (2H, s), 4.77 (2H, s), 6.94 (1H, d), 7.29 (1H, dd), 7.48 (2H, m), 7.56 (1H, d), 7.77-7.89 (4H, m), 8.23 (1H, d)

Compound 64:

45 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.18 (3H, s), 3.59 (3H, s), 3.83 (3H, s), 4.26 (2H, s), 4.77 (2H, s), 4.82 (2H, s), 6.91 (1H, d), 7.11 (1H, dd), 7.26-7.58 (7H, m), 7.72-7.88 (5H, m), 8.20 (1H, d)

Compound 65:

50 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.25 (3H, s), 3.20 (3H, s), 3.76 (3H, s), 3.84 (3H, s), 5.07 (2H, s), 6.93 (1H, d), 7.24 (1H, dd), 7.44-7.55 (4H, m), 7.76-7.89 (3H, m), 7.98 (1H, d)

Compound 66:

55 $^1\text{H-NMR}(\text{CDCl}_3)$ δ ppm: 2.20 (3H, s), 2.43 (3H, s), 3.55 (3H, s), 3.74 (3H, s), 5.08 (2H, s), 6.85 (1H, d), 7.23 (1H, dd), 7.30 (2H, dd), 7.41-7.52 (4H, m), 7.55-7.88 (6H, m)

Compound 67:

¹H-NMR(CDCl₃) δ ppm: 1.24 (3H, s), 2.62 (3H, s), 2.98 (3H, s), 3.84 (3H, s), 4.71 (2H, bs), 6.99 (1H, d), 7.30 (1H, dd), 7.38-7.52 (4H, m), 7.58 (1H, d), 7.87 (2H, t), 8.17 (1H, d)

Compound 68:

¹H-NMR(CDCl₃) δ ppm: 1.16 (3H, t), 1.24 (3H, t), 2.54 (2H, q), 2.62 (2H, q), 3.80 (3H, s), 4.9-5.1 (2H, b), 6.91 (1H, d), 7.26 (1H, dd), 7.45 (2H, m), 7.54 (1H, d), 7.77 (1H, d), 7.84 (1H, d), 7.87 (1H, d), 8.01 (1H, d)

Compound 69:

¹H-NMR(CDCl₃) δ ppm: 0.88 (3H, t), 1.24 (3H, t), 1.65 (2H, m), 2.49 (2H, t), 2.63 (2H, q), 3.79 (3H, s), 4.9-5.1 (2H, b), 6.91 (1H, d), 7.26 (1H, dd), 7.45 (2H, m), 7.54 (1H, d), 7.77 (1H, d), 7.84 (1H, d), 7.87 (1H, d), 8.01 (1H, d)

Compound 70:

¹H-NMR(CDCl₃) δ ppm: 0.97 (3H, t), 2.33 (2H, q), 3.74 (3H, s), 5.17 (2H, bs), 6.94 (1H, d), 7.21-7.38 (4H, m), 7.4-7.5 (4H, m), 7.56 (1H, d), 7.77 (1H, dd), 7.84 (1H, d), 7.86 (1H, d), 7.96 (1H, dd), 8.39 (1H, d)

Compound 71:

¹H-NMR(CDCl₃) δ ppm: 1.21 (3H, t), 2.44 (2H, q), 2.58 (2H, q), 3.59 (3H, s), 5.05 (2H, bs), 6.92 (1H, d), 7.25-7.34 (3H, m), 7.47 (2H, m), 7.4-7.5 (2H, m), 7.68 (2H, d), 7.79 (1H, d), 7.85 (1H, d), 7.88 (1H, d), 7.96 (1H, d), 7.98 (1H, dd), 8.39 (1H, d)

Formulation Examples of the agricultural and horticultural fungicides, bacteriocides, insecticides, and acaricides containing the compound of the present invention as an active ingredient are shown below for illustrative purposes only but not for limitation.

FORMULATION EXAMPLE 1

Wettable Powder:

Twenty parts of the compound of the present invention, 20 parts of Carplex #80 (white carbon produced by Shionogi & Co., Ltd.), 52 parts of ST Kaolin Clay (kaolinite produced by Tsuchiya Kaolin K.K.), 5 parts of Solpol 9047K (an anionic surface active agent produced by Toho Chemical Industry Co., Ltd.), and 3 parts of Runox P65L (an anionic surface active agent produced by Toho Chemical Industry Co., Ltd.) were mixed and ground uniformly to obtain a wettable powder containing 20% of the active ingredient.

FORMULATION EXAMPLE 2

Dust:

Two parts of the compound of the present invention, 93 parts of clay (produced by Nippon Talc K.K.), and 5 parts of Carplex #80 (white carbon produced by Shionogi & Co., Ltd.) were uniformly mixed and ground to obtain a dust containing 2% of the active ingredient.

FORMULATION EXAMPLE 3

Emulsifiable Concentrate:

In a mixed solvent of 35 parts of xylene and 30 parts of dimethylformamide was dissolved 20 parts of the compound of the present invention, and 15 parts of Solpol 3005X (a mixture of a nonionic surface active agent and an anionic surface active agent, produced by Toho Chemical Co., Ltd.) was added thereto to prepare an emulsifiable concentrate containing 20% of the active ingredient.

FORMULATION EXAMPLE 4

Flowable:

5 A mixture of 30 parts of the compound of the present invention, 5 parts of Solpol 9047K, 3 parts of Sorbon T-20 (a nonionic surface active agent produced by Toho Chemical Co., Ltd.), 8 parts of ethylene glycol, and 44 parts of water were wet ground in Dynamill (produced by shinmaru enterprises Co.). To the resulting slurry was added 10 parts of a 1% aqueous solution of xanthan gum (naturally occurring polymer), followed by mixing and grinding thoroughly to obtain a flowable containing 20% of the active ingredient.

TEST EXAMPLE 1Insecticidal Effect on Larvae of *Nilaparvata lugens*:

15 A rice seedling was setted in a glass cylinder (inner diameter: 3 cm; length: 17 cm), and five 4th instar larvae of *Nilaparvata lugens* were set free therein. The agricultural and horticultural pesticide of the present invention (an emulsifiable concentrate) was prepared in accordance with Formulation Example 3 and diluted with water, and 0.5 ml of the resulting emulsion was sprayed in the cylinder by means of a spray tower (manufactured by Mizuho Rika) (duplicates at 1 concentration). Five days after the treatment, the mortality and agony of the larvae were examined to obtain a death rate (%) taking an agonizing insect as a 1/2 dead insect. The results obtained are shown in Table 4 (the compound numbers in Table 4 correspond to those in Tables 1, 2 and 3).

TABLE 4

Compound No.	Concentration (ppm)	Death Rate (%)
1	500	100
2	500	100
7	500	100
8	500	100
9	500	100
10	500	100
11	500	100
12	500	100
13	500	100
18	500	100
25	500	90
26	500	90
27	500	100
30	500	100
31	500	90
34	500	100
35	500	100

TABLE 4 (cont'd.)

Compound No.	Concentration (ppm)	Death Rate (%)
38	500	90
40	500	100
42	500	90
43	500	90
50	500	100
51	500	100
52	500	90
53	500	90
54	500	100
55	500	100
56	500	100
57	500	100
63	500	100

TEST EXAMPLE 2

Insecticidal Effect on Larvae of *Plutella xylostella*:

The agricultural and horticultural pesticide of the present invention (a wettable powder) was obtained in accordance with Formulation Example 1 and diluted with water to prepare a suspension having a prescribed concentration. A disc (6 cm in diameter) cut out of a cabbage leaf was soaked in the suspension for 1 minute, air-dried, and placed in a plastic cup (inner diameter: 7 cm). Five 3-instar larvae of *Plutella xylostella* were put in the cup (duplicates at one concentration). Four days later, the death and agony of the larvae were examined to obtain a death rate (%) taking an agonizing insect as a 1/2 dead insect. The results obtained are shown in Table 5 (the compound numbers in Table 5 correspond to those in Tables 1, 2 and 3).

TABLE 5

Compound No.	Concentration (ppm)	Death Rate (%)
4	500	100
5	500	100
7	500	100
10	500	100
11	500	100
12	500	100
13	500	100
18	500	100
25	500	90
27	500	90
28	500	100
31	500	100
34	500	100
38	500	90
39	500	90
40	500	100
42	500	100
43	500	100
51	500	100
62	500	100

TEST EXAMPLE 3

Acaricidal Effect on Imagoes of *Tetranychus urticae*:

Ten female imagoes of *Tetranychus urticae* were set free on a disc (3 cm in diameter) cut out of a kidney bean leaf. The agricultural and horticultural pesticide of the present invention (a wettable powder) was prepared in accordance with Formulation Example 1 and diluted with water to a prescribed concentration, and 3.5 ml of the resulting suspension was sprayed on the leaf disc by means of a rotary spray tower (manufactured by Mizuho Rika) (duplicates at one concentration). Twenty-four hours later dead mites were counted to obtain an imago death rate (%). The results obtained are shown in Table 6.

TEST EXAMPLE 4

Acaricidal Effect on Eggs of *Tetranychus urticae*:

5 Five female imagoes of *Tetranychus urticae* were set free on a disc (3 cm in diameter) of a kidney bean leaf for 20 hours to let the mites lay eggs and then removed. The agricultural and horticultural pesticide of the present invention (a wettable powder) was prepared in accordance with Formulation Example 1 and diluted with water to a prescribed concentration, and 3.5 ml of the resulting suspension was sprayed on the leaf disc by means of a rotary spray tower (manufactured by Mizuho Rika) (duplicates at one concentration). After 8 days from the application, hatched eggs and
10 unhatched eggs were counted to obtain an egg death rate (%). The results obtained are shown in Table 6.

TABLE 6

Compound No.	Concentration (ppm)	Imago Death Rate (%)	Egg Death Rate (%)
2	500	100	100
12	500	100	100

TEST EXAMPLE 5

Fungicidal Effect on Rice Blast Fungus (*Magnaporthe grisea*):

Ten/pot rice plants (variety: Akinishiki) were cultivated in a resin pot (diameter: 6 cm). The agricultural and horticultural pesticide of the present invention (a wettable powder) was prepared in accordance with Formulation Example 1 and diluted with water to a prescribed concentration. The foliage of the plants at 3 to 4 leaf stage was sprayed with 10 ml/pot of the diluted preparation. After air drying, the plants were spray-inoculated with a spore suspension (5.0×10^5 spores/ml) of *Magnaporthe grisea* having been cultured on an oatmeal medium, placed in a moist chamber at 25°C for 24 hours, and then allowed to stand in a greenhouse (22 to 25°C) for 7 days. The number of lesions appearing on the leaves were counted to obtain an average per leaf, and a control value was calculated according to the following formula. The results are shown in Table 7.

$$\text{Control value (\%)} = (\text{number of lesions per leaf in untreated pot} - \text{number of lesions per leaf in treated pot}) / (\text{number of lesions per leaf in untreated pot}) \times 100$$

TABLE 7

Compound No.	Concentration (ppm)	Control Value (%)
3	250	100
9	250	91
10	250	89
11	250	85
12	250	100
13	250	87
24	250	100
27	250	90
34	250	90
35	250	96
38	250	100
39	250	100
40	250	100
45	250	90
46	250	96
47	250	92
48	250	100
49	250	100
50	250	97
54	250	100
55	250	100
56	250	97
62	250	100
63	250	98

TEST EXAMPLE 6

Fungicidal Effect on Tomato Late Blight Fungus (*Phytophthora infestans*):

Three/pot tomato plants (variety: Red Cherry) were cultivated in a resin pot (diameter: 6 cm). The agricultural and horticultural pesticide of the present invention (a wettable powder) was prepared in accordance with Formulation Example 1 and diluted with water to a prescribed concentration. The resulting suspension was sprayed to the foliage of the plants at 3 to 4 leaf stage in an amount of 10 ml per pot. After air drying the suspension, the plants were spray-inocu-

EP 0 891 975 A1

lated with a suspension of zoosporendia (5.9×10^4 zoosporendia/ml) of *Phytophthora infestans* having been cultured on tomato leaves cut off a tomato plant, and the inoculated plants were placed in a moist chamber at 25°C for 24 hours and then allowed to stand in a greenhouse (20 to 22°C) for 4 days. The total area of the lesions appearing on each leaf was measured and expressed in disease severity index. A degree of disease (%) was calculated from the disease severity indices, and a control value (%) was obtained therefrom. The results are shown in Table 8. Disease severity index:

- 0 No lesion.
- 1 Lesion area is 1/3 or less.
- 3 Lesion area is 1/3 to 2/3.
- 5 Lesion area is 2/3 or more.

$$\text{Degree of disease (\%)} = (0 \times n_0 + 1 \times n_1 + 3 \times n_3 + 5 \times n_5) / (n_0 + n_1 + n_3 + n_5) \times 100$$

wherein n_x is the number of leaves having a disease severity index of x .

$$\text{Control value (\%)} = (\text{degree of disease in untreated pot} - \text{degree of disease in treated pot}) / \text{degree of disease in untreated pot} \times 100$$

TABLE 8

Compound No.	Concn. (ppm)	Control Value (%)
3	250	100
4	250	94
5	250	100
6	250	91
7	250	95
8	250	100
9	250	100
10	250	100
11	250	92
12	250	89
13	250	85
18	250	100
22	250	100
24	250	100
26	250	100
27	250	100
28	250	100
32	250	100
34	250	100
35	250	93
39	250	100
40	250	100
41	250	100
42	250	100
43	250	100

TABLE 8 (cont'd)

Compound No.	Concn. (ppm)	Control Value (%)
46	250	97
48	250	100
52	250	100
54	250	96
55	250	92
56	250	90
62	250	100
64	250	96
66	250	99

TEST EXAMPLE 7

Acute Toxicity in Mice in Single Oral Administration:

Acute toxicity (p.o.) in mice of compound Nos. 8, 10, 13, 22, 23, 38, 42, and 52 was examined as follows. For comparison, compounds (IV), (V) and (VI) disclosed in JP-A-2-62876 were tested similarly.

Compound Nos. 8, 10, 22, 23, 38, and 42 (30 mg each) were each suspended in 10 ml of a 0.5% CMC-Na aqueous solution. Compound Nos. 13 and 52 and compounds (IV), (V) and (VI) (30 mg each) were each suspended in 10 ml of a 0.5% tragacanth gum aqueous solution. The suspension was orally given to a group each consisting of five 6-week-old CD-1 male mice (available from Charles River) at a dose of 10 ml-suspension per kg of the body weight. The number of dead mice was counted after 7 days from the administration to obtain a death rate. The results obtained are shown in Table 9 below.

TABLE 9


Compound No.	Dose (mg-active ingredient/kg)	Death Rate (%)
8	30	0
10	30	0
13	30	0
22	30	0
23	30	0
38	30	0
42	30	0
52	30	0
(IV)	30	100
(V)	30	100
(VI)	30	100

Industrial Applicability:

The pyrazole compounds of the present invention exhibit excellent controlling effects on various plant pathogenic fungi or bacteria, insect pests and mites and are of high safety. The pyrazole compounds of the present invention are very useful as an active ingredient of fungicides, bacteriocides, insecticides and acaricides.

Claims


1. A pyrazole compound represented by formula (I):



wherein R^1 represents a hydrogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a phenyl group; R^2 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_{1-4} alkyl group, a C_{1-4} alkoxy group, a C_{1-4} haloalkoxy group, a C_{2-8} alkoxyalkoxy group or a benzyloxy group; or R_1 and R_2 are taken together to form a group:




wherein R^4 represents a hydrogen atom or a C_{1-4} alkyl group;
 R^3 represents a hydrogen atom, a C_{1-4} alkyl group, a C_{2-5} alkenyl group, a C_{5-7} cycloalkenyl group, a C_{2-5} alkoxyalkyl group, a C_{8-12} benzyloxyalkyl group, a formyl group, a C_{2-5} alkanoyl group, a benzoyl group, a C_{2-5} alkoxycarbonyl group, a benzyloxycarbonyl group, a C_{1-4} alkylsulfonyl group or a C_{6-10} arylsulfonyl group;
 and R represents a group represented by formulae:



wherein R^5 represents a hydrogen atom, a halogen atom or a C_{1-4} alkoxy group; R^6 represents a halogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a C_{1-4} haloalkoxy group; and n represents 1 or 2.

2. The pyrazole compound according to claim 1, wherein, in the general formula (I) shown in claim 1, R is a group represented by formula:




wherein R^5 represents a hydrogen atom, a halogen atom or a C_{1-4} alkoxy group; and n represents 1 or 2.

3. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R¹ is a hydrogen atom or a C₁₋₄ alkyl group.
4. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R² is a halogen atom, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group.
5. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R₁ and R₂ are taken together to form a group represented by formula:



wherein R⁴ represents a hydrogen atom or a C₁₋₄ alkyl group.

6. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R³ is a hydrogen atom, a C₂₋₅ alkenyl group, a C₅₋₇ cycloalkenyl group, a C₂₋₅ alkoxyalkyl group, a benzyloxymethyl group, a formyl group, a C₂₋₅ alkanoyl group or a p-toluenesulfonyl group.
7. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R³ is a hydrogen atom, a formyl group or an alkanoyl group.
8. The pyrazole compound according to claim 1, wherein, in the general formula (I) shown in claim 1, R is a group represented by formula:



wherein R⁶ is a halogen atom, a C₁₋₄ alkyl group, a C₁₋₄ haloalkyl group, a C₁₋₄ alkoxy group or a C₁₋₄ haloalkoxy group.

9. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R¹ is a C₁₋₄ alkyl group or a C₁₋₄ haloalkyl group.
10. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R² is a hydrogen atom, a halogen atom, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group.
11. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R₁ and R₂ are taken together to form a group represented by formula:



wherein R⁴ represents a hydrogen atom or a C₁₋₄ alkyl group.

12. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R³ is a hydrogen atom, a C₂₋₅ alkenyl group, a C₅₋₇ cycloalkenyl group, a C₂₋₅ alkoxyalkyl group, a benzyloxymethyl group, a formyl group, a C₂₋₅ alkanoyl group or a p-toluenesulfonyl group.

13. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 2, R^3 is a hydrogen atom, a formyl group or an alkanoyl group.
14. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R^6 is a halogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a C_{1-4} haloalkoxy group.
15. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R^6 is a halogen atom or a C_{1-4} haloalkyl group.
16. A pesticide comprising the pyrazole compound according to claim 1.
17. A fungicide or bactericide comprising the pyrazole compound according to claim 1.
18. An insecticide comprising the pyrazole compound according to claim 1.
19. An acaricide comprising the pyrazole compound according to claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/01162

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl⁶ C07D401/12, 14, 405/14, 417/14, A01N43/56, 78

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int. Cl⁶ C07D401/12, 14, 405/14, 417/14, A01N43/56, 78

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP, 2-62876, A (Mitsubishi Chemical Corp.), March 2, 1990 (02. 03. 90), Full descriptions & EP, 329090, A & US, 4968805, A	1, 8-12, 14-19
A		2-7, 13
P, A	JP, 8-277287, A (Mitsubishi Chemical Corp.), October 22, 1996 (22. 10. 96), Claim & EP, 726266, A & AU, 9643311, A	1 - 19

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

June 16, 1997 (16. 06. 97)

Date of mailing of the international search report

June 24, 1997 (24. 06. 97)

Name and mailing address of the ISA/

Japanese Patent Office

Facsimile No.

Authorized officer

Telephone No.



European Patent
Office


**SUPPLEMENTARY
EUROPEAN SEARCH REPORT**

Application Number
EP 97 91 4601

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
	No further relevant documents disclosed -----		C07D401/12 C07D401/14 C07D405/14 C07D417/14 A01N43/56 A01N43/78
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07D
The supplementary search report has been based on the last set of claims valid and available at the start of the search.			
Place of search		Date of completion of the search	Examiner
MUNICH		24 January 2000	Fink, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			


WHAT IS CLAIMED IS:

1. A pyrazole compound represented by formula (I):




(I)

wherein R¹ represents a hydrogen atom, a C₁₋₄ alkyl group, a C₁₋₄ haloalkyl group, a C₁₋₄ alkoxy group or a phenyl group; R² represents a hydrogen atom, a halogen atom, a hydroxyl group, a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, a C₁₋₄ haloalkoxy group, a C₂₋₈ alkoxyalkoxy group or a benzyloxy group; or R₁ and R₂ are taken together to form a group:



wherein R⁴ represents a hydrogen atom or a C₁₋₄ alkyl group;

R³ represents a hydrogen atom, a C₁₋₄ alkyl group, a C₂₋₅ alkenyl group, a C₃₋₇ cycloalkenyl group, a C₂₋₅ alkoxyalkyl group, a C₈₋₁₂ benzyloxyalkyl group, a formyl group, a C₂₋₅ alkanoyl group, a benzoyl group, a C₂₋₅ alkoxycarbonyl group, a benzyloxycarbonyl group, a C₁₋₄ alkylsulfonyl group or a C₆₋₁₀ arylsulfonyl group; and R represents a group represented by formulae:




5

wherein R^5 represents a hydrogen atom, a halogen atom or a C_{1-4} alkoxy group; R^6 represents a halogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a C_{1-4} haloalkoxy group; and n represents 1 or 2.

10

2. The pyrazole compound according to claim 1, wherein, in the general formula (I) shown in claim 1, R is a group represented by formula:

15



wherein R^5 represents a hydrogen atom, a halogen atom or a C_{1-4} alkoxy group; and n represents 1 or 2. ✓

3. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R^1 is a hydrogen atom or a C_{1-4} alkyl group. ✓

20

4. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R^2 is a halogen atom, a C_{1-4} alkyl group or a C_{1-4} alkoxy group. ✓

5. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R₁ and R₂ are taken together to form a group represented by formula:




5 wherein R⁴ represents a hydrogen atom or a C₁₋₄ alkyl group. ✓

6. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R³ is a hydrogen atom, a C₂₋₅ alkenyl group, a C₅₋₇ cycloalkenyl group, a C₂₋₅ alkoxyalkyl group, a benzyloxymethyl group, a formyl group, a C₂₋₅ alkanoyl group or a p-toluenesulfonyl group.

7. The pyrazole compound according to claim 2, wherein, in the general formula (I) shown in claim 2, R³ is a hydrogen atom, a formyl group or an alkanoyl group. ✓

8. The pyrazole compound according to claim 1, wherein, in the general formula (I) shown in claim 1, R is a group represented by formula:



20 wherein R⁶ is a halogen atom, a C₁₋₄ alkyl group, a C₁₋₄ haloalkyl group, a C₁₋₄ alkoxy group or a C₁₋₄ haloalkoxy group. ✓

9. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R¹ is a C₁₋₄ alkyl group or a C₁₋₄ haloalkyl group. ✓

10. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R^2 is a hydrogen atom, a halogen atom, a C_{1-4} alkyl group or a C_{1-4} alkoxy group. ✓

5 11. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R_1 and R_2 are taken together to form a group represented by formula:



10 wherein R^4 represents a hydrogen atom or a C_{1-4} alkyl group. ✓

12. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R^3 is a hydrogen atom, a C_{2-5} alkenyl group, a C_{5-7} cycloalkenyl group, a C_{2-5} alkoxyalkyl group, a benzyloxymethyl group, a formyl group, a C_{2-5} alkanoyl group or a p-toluenesulfonyl group. ✓

15 13. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 2, R^3 is a hydrogen atom, a formyl group or an alkanoyl group. ✓

20 14. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R^6 is a halogen atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, a C_{1-4} alkoxy group or a C_{1-4} haloalkoxy group. ✓

25 15. The pyrazole compound according to claim 8, wherein, in the general formula (I) shown in claim 8, R^6 is a halogen atom or a C_{1-4} haloalkyl group. ✓

16. A pesticide comprising the pyrazole compound according to claim 1.

17. A fungicide or bacteriocide comprising the pyrazole compound according to claim 1.

5 18. An insecticide comprising the pyrazole compound according to claim 1.

19. An acaricide comprising the pyrazole compound according to claim 1.